

---

# MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2019

January–March 2019—Volume 134, pp. 161–167

<https://doi.org/10.5248/134.161>

---

## ***Spadicoides matsushimae* sp. nov., and *Anisospadicoides* gen. nov. for two atypical *Spadicoides* species**

MIN QIAO<sup>1</sup>, DE-WEI LI<sup>2,3</sup>, ZE-FEN YU<sup>1\*</sup>,  
KAI ZHANG<sup>4</sup>, RAFAEL F. CASTAÑEDA-RUIZ<sup>5</sup>

<sup>1</sup> *Laboratory for Conservation and Utilization of Bio-resources,  
Key Laboratory for Microbial Resources of the Ministry of Education,  
Yunnan University, Kunming, Yunnan, 650091, China*

<sup>2</sup> *The Connecticut Agricultural Experiment Station,  
Valley Laboratory, 153 Cook Hill Road, Windsor, CT 06095, USA*

<sup>3</sup> *Co-Innovation Center for Sustainable Forestry in Southern China,  
Nanjing Forestry University, Nanjing, Jiangsu 210037, China*

<sup>4</sup> *Department of Landscaping, Shandong Yingcai University, Jinan, 250104, China*

<sup>5</sup> *Instituto de Investigaciones Fundamentales en Agricultura Tropical Alejandro de Humboldt  
(INIFAT), Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200*

\*CORRESPONDENCE TO: [zfyuqm@hotmail.com](mailto:zfyuqm@hotmail.com)

**ABSTRACT**—*Anisospadicoides* is proposed as a new genus for two species of *Spadicoides*, *S. macrocontinua* and *S. macroobovata*, that have polytretic and blastic conidial ontogeny on the apical conidiogenous cells. A new species, *Spadicoides matsushimae*, distinguished by broadly fusiform to navicular, mostly 2-septate, brown to dark brown conidia, is also described and illustrated.

**KEY WORDS**—asexual *Ascomycota*, *Helminthosphaeriaceae*, hyphomycetes, systematics

### **Introduction**

*Spadicoides* S. Hughes, typified by *S. bina* (Corda) S. Hughes, is distinguished by macronematous, mononematous, and generally unbranched erect conidiophores. Conidiogenous cells are polytretic, integrated, terminal and intercalary, determinate, with conidiogenous loci

at the apex and laterally below. The conidia are solitary, variously shaped (ellipsoidal, oblong, obovoid, obclavate, globose, clavate, or cuneate), unicellular or multicellular, euseptate, smooth or verruculose, subhyaline to brown or black, and dry (Ellis 1971, Hughes 1958, Ma & al. 2016, Xia & al. 2013, Whitton & al. 2012).

Two *Spadicoides* species, *S. macrocontinua* and *S. macrobovata*, have an unusual combination of tretic and holoblastic conidial development on the apical conidiogenous cells (Matsushima 1993, 1995). Conidial ontogeny and conidiogenous events have been considered the fundamental criteria for asexual fungal taxonomy and generic delimitation (Kendrick 2017, Seifert & al. 2011). Consequently, we propose the new genus *Anisospadicoides* to accommodate two atypical *Spadicoides* species.

In 1993, Matsushima published a description and illustrations of a conspicuous fungus found on rotten petiole of a palm, but he did not provide a species epithet for it, merely calling it “*Spadicoides* sp.” This fungus is formally named here as a new species, *S. matsushimae*.

## Taxonomy

*Anisospadicoides* R.F. Castañeda, Qiao & Z.F. Yu, gen. nov.

FIG. 1

MYCOBANK MB 827953

Differs from *Spadicoides* species by its holoblastic, mostly monoblastic conidial ontogeny, and production of conidia on the apical conidiogenous cells beside or near the tretic loci.



TYPE SPECIES: *Spadicoides macrocontinua* Matsush. [= *Anisospadicoides macrocontinua* (Matsush.) R.F. Castañeda & al.]

ETYMOLOGY: Greek, *aniso-* meaning unequal, uneven, or dissimilar + Latin, *-spadicoides*, referring to the genus *Spadicoides*.

COLONIES on CMA medium felted or hairy, brown to dark brown. CONIDIOPHORES macronematous, mononematous, unbranched, erect, straight, cylindrical, septate, smooth, brown to dark brown. CONIDIOGENOUS CELLS biontogenous, sometimes on the same cell: i) polytretic, integrated, mostly intercalary, cylindrical, sometimes terminal with subapical tretic loci and apical blastic loci; ii) monoblastic, integrated, terminal, sometimes with several tretic loci near below the blastic locus on the same cell.

---

FIG. 1. *Anisospadicoides macrocontinua* (holotype, MFC OP-302): A. Conidiogenous cells with apical conidia originated by holoblastic mode and subapical conidia borne from tretic loci; B. Conidiogenous cells with apical conidia originated by holoblastic mode and subapical tretic loci.






FIG. 2. *Anisospadicoides macrobovata* (holotype, MFC 4P-562): A. Conidia produced after holoblastic mode on the apical loci; B. Conidia produced after enteroblastic mode on the subapical and intercalary loci; C. Conidiogenous cells with apical blastic loci and subapical and intercalary tretic loci.

CONIDIA solitary, dimorphic: i) obovoid, subpyriform, truncate at the base, acropleurogenous, mostly pleurogenous, unicellular or septate, smooth, brown to dark brown or black, dry, borne on tretic conidiogenous loci; ii) obovoid, broadly subfusiform, to subclavate, attenuate or slightly fimbriate towards the truncate base, acrogenous, unicellular or septate, euseptate, smooth, brown to dark brown or black, dry, borne on blastic conidiogenous loci.

*Anisospadicoides macrocontinua* (Matsush.) R.F. Castañeda, Qiao & Z.F. Yu, **comb. nov.** FIG. 1  
MYCOBANK MB 827954  
= *Spadicoides macrocontinua* Matsush., Matsush. Mycol. Mem.7: 67 (1993).

*Anisospadicoides macroobovata* (Matsush.) Qiao, Z.F. Yu & R.F. Castañeda, **comb. nov.** FIG. 2  
MYCOBANK MB 827955  
= *Spadicoides macroobovata* Matsush., Matsush. Mycol. Mem. 8: 36 (1995).

*Spadicoides matsushimae* R.F. Castañeda & D.W. Li, **sp. nov.** FIG. 3  
MYCOBANK MB 809530

Differs from *Spadicoides curvularioides* by its broadly fusiform, smaller, 2-septate, brown conidia.

TYPE: Peru, Rio Sinchicuy, 14°36'N 90°38'W, on decaying petiole of an unidentified palm, June 1991, coll. T. Matsushima (**Holotype**, MFC 1P-294).

ETYMOLOGY: Latin, *matsushimae*, in reference to Takashi Matsushima, collector of the holotype specimen.

CONIDIOPHORES on the natural substratum macronematous, mononematous, unbranched, erect, straight, rigid, septate, brown, smooth, 50–160 µm long, 4.5–6 µm wide at the base, CONIDIOGENOUS CELLS polytretic, integrated, terminal and intercalary, determinate, brown. CONIDIA solitary, acropleurogenous, broadly fusiform to broadly navicular, conical-truncate at the base, rounded at the apex, (1–) 2-septate, smooth, 15–19 × 7–9 µm, with central cell brown to dark brown and larger than pale brown end cells.

NOTES: There are 58 previously described species of *Spadicoides* (Index Fungorum 2018), of which only 42 were accepted by Goh & Hyde (1996), Kuthubutheen & Nawawi (1991), Ma & al. (2014, 2016), Whitton & al. (2012), and Xia & al. (2013). Ma & al. (2016) provided a comparative table of *Spadicoides* spp. and a key for their identification. Among the recognized *Spadicoides* species, only *S. curvularioides* B. Sutton & Hodges is superficially similar to *S. matsushimae* in producing more or less fusiform or navicular




FIG. 3. *Spadicoides matsushimae* (holotype, MFC 1P-294):  
A. Conidia; B. Conidiogenous cells and conidiogenous loci.

conidia, but *S. curvularioides* has pale brown conidia that are paler at the ends, 3-septate, longer (22–33.5 µm), and verruculose (Sutton & Hodges 1978).

### Acknowledgments

This work was financed by the National Natural Science Foundation Program of PR China (31760012, 31570023). The authors express their sincere gratitude to Dr. Xiu-Guo Zhang and Dr. Josiane S. Monteiro for their critical review of the manuscript. We acknowledge the facilities provided by Dr. P.M. Kirk and Dr. K. Bensch through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

### Literature cited

- Ellis MB. 1971. Dematiaceous hyphomycetes. Kew, Commonwealth Mycological Institute.
- Goh TK, Hyde KD. 1996. *Spadicoides cordanoides* sp. nov., a new dematiaceous hyphomycete from submerged wood in Australia, with a taxonomic review of the genus. *Mycologia* 88(6): 1022–1031. <https://doi.org/10.1080/00275514.1996.12026745>
- Hughes SJ. 1958. Revisiones hyphomycetum aliquot cum appendice de nominibus rejiciendis. *Canadian Journal of Botany* 36: 727–836. <https://doi.org/10.1139/b58-067>
- Index Fungorum. 2018. [www.indexfungorum.org/Names/Names.asp](http://www.indexfungorum.org/Names/Names.asp) (accessed 29 Sept. 2018).
- Kendrick B. 2017. *The fifth kingdom* (4<sup>th</sup> ed). Hackett publishing company. 416 p.
- Kuthubutheen AJ, Nawawi A. 1991. Two new species of *Spadicoides* from Malaysia. *Mycological Research* 95: 163–168. [https://doi.org/10.1016/S0953-7562\(09\)81006-X](https://doi.org/10.1016/S0953-7562(09)81006-X)
- Ma LG, Xia JW, Ma YR, Castañeda-Ruiz RF, Zhang XG. 2014. Two new species of *Spadicoides* and *Gangliostilbe* from southern China. *Mycological Progress* 13: 547–552. <https://doi.org/10.1007/s11557-013-0937-z>
- Ma J, Castañeda-Ruiz RF, Zhang XG. 2016. Three new species of *Spadicoides* from Lushan Mountain. *Mycological Progress* 15: 43 [8 p.]. <https://doi.org/10.1007/s11557-016-1185-9>
- Matsushima T. 1993. *Matsushima Mycological Memoirs* no. 7. Matsushima Fungus Collection, Kobe, Japan.
- Matsushima T. 1995. *Matsushima Mycological Memoirs* no. 8. Matsushima Fungus Collection, Kobe, Japan.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. *CBS Biodiversity Series* 9. 997 p.
- Sutton BC, Hodges CS. 1978. *Eucalyptus* microfungi: *Chaetendophragmiopsis* gen. nov. and other hyphomycetes. *Nova Hedwigia* 29: 593–607.
- Whitton SR, McKenzie EHC, Hyde KD. 2012. Anamorphic fungi associated with *Pandanaceae*. 125–353, in: SR Whitton & al. (eds). *Fungi Associated with Pandanaceae*. *Fungal Diversity Research Series* 21. [https://doi.org/10.1007/978-94-007-4447-9\\_4](https://doi.org/10.1007/978-94-007-4447-9_4)
- Xia JW, Ma LG, Ma J, Zhang XG. 2013. Two new species of *Spadicoides* from southern China. *Mycotaxon* 126: 55–60. <https://dx.doi.org/10.5248/126.55>